

Archives of Psychiatry and Mental Health

Volume - 6, Issue - 1

Mini Review

Published Date:- 2022-12-23

[The necessity of emotion management for the general public: a comparison of diagnostic changes in two anger-related psychiatric disorders](#)

Although emotion management is essential to mental health, public health has not paid much attention to emotional conditions or disorders. By analyzing a nine-year diagnostic trend and sociodemographic characteristics of two mental disorders: Hwabyung and Intermittent Explosive Disorder (IED), this study demonstrated the importance of managing emotions precautionarily.

Data reconstructed from the National Health Insurance Service were used to analyze the yearly diagnostic trend in the two mental disorders characterized by anger.

Hwabyung was more common among women and middle-aged individuals, despite the varied number of diagnoses by year. Between 2010 and 2018, IED diagnoses gradually rose, with the average rate of increase the highest in the 20s for male IED diagnoses in 2017. The low prevalence of the IED in Korea compared to other Western countries and the gender and age differences in both Hwabyung and IED diagnoses suggest the role of cultural influences related to emotions (or emotional management).

Especially in light of the world's emphasis on resilience to COVID-19, these results indicate how public emotional management is essential during stressful situations. The results also highlight the need for community mental health programs tailored to gender and age.

Research Article

Published Date:- 2022-11-30

[The role of denial on emotional correlates of childhood trauma in a non-clinical population](#)

Background: A large body of evidence suggests that child abuse and neglect by a caregiver is a recurrent event linked to increased psychopathology symptoms. The Childhood Trauma Questionnaire (CTQ) is commonly used to assess abuse/neglect during childhood. However, even though the Minimization-Denial (MD) subscale was originally designed to assess response bias (i.e., underreporting of childhood maltreatment), it is possible that the scale may reflect coping strategies that play an effective role in the relationship between childhood trauma and their negative outcomes. Also, even though MD has been associated with decreased psychopathology symptoms, it is also strongly associated with other scales of the CTQ.

Method: This study (n = 133) examined whether (1) the MD-scale is negatively associated with alexithymia, emotion dysregulation and psychopathology, if (2) these associations will hold when adjusting for different subtypes of abuse and neglect and (3) and the role of MD as a possible moderator in these relationships.

Results: The analyses showed that, although MD scores have relatively strong and (mostly) significant (negative) associations with the CTQ, emotion dysregulation strategies and psychopathology symptoms, these associations were weak and failed to remain significant when adjusting for the effect of CTQ.

Conclusion: Our findings suggest that the MD scores should be viewed as an accurate reflection of the absence (or little) of exposure to childhood abuse/neglect.

Review Article

Published Date:- 2022-10-26

[The central role of desire in mediating bullying behavior in schools](#)

Numerous studies of bullying in schools have identified aspects of the environment (E) and aspects of the person (P) as contributing to the prevalence of bullying in schools. It is proposed that the concept of desire can play a central role in explaining how bullying in schools arises and how it can be effectively addressed by schools by promoting social and emotional learning, mindfulness, and problem-solving techniques such as the Method of Shared Concern. The article identifies a need for further research to confirm the hypothesized relationships and assess the utility of the concept of desire as a variable in research and the practice of bullying prevention in schools.

[Being a parent of a child with cancer: What psychosocial and family repercussions](#)

Cancer is a serious disease that affects deeply and painfully not only the child who has cancer but also their parents. Through this study, we describe the different aspects of the impact of pediatric cancer on parents: the psychological, social, and family impact to offer optimal care to these parents.

Results: mothers represent 82.5% of the participants in our survey. More than sixty percent were of urban origin. The average time from diagnosis to parents' assessment was 7.3 months. This announcement was made by doctors in 87.5% of cases. Conscious denial of cancer when it was announced was reported in 75% of parents. The social impact of pediatric cancer on parents was significant. The child's illness was experienced as a very significant psychological distress; all of the parents said they had given up on important projects after their children's illness. The psycho-emotional impact was represented by feelings of guilt in 37.5% and incapacity for illness in 30%. Forty-two percent felt tensions on the marital level with significant repercussions on the couple with a type of destabilization in 60% of cases. The parent's relationship with the rest of the family, especially siblings, was marked by neglect and anxiety in 35% and 26% respectively.

Conclusion: The discovery of pediatric cancer induces various feelings that will inevitably have an impact on the parents of the affected child. Understanding the different aspects of this impact on the parents' psycho-social, emotional and family experiences will make it possible to offer optimal care.

Case Report**Published Date:- 2022-06-15**[Prevalence of PTSD symptoms in children: The cost of the inadequate mental health system in Pakistan](#)

Post-Traumatic Stress Disorder (PTSD) is a disorder that affects not only military veterans but also the general public specifically children. They may replay the event and its aftermath, avoid talking about it, have low self-esteem, and be unable to develop meaningful connections. Mentally sick individuals in Pakistan frequently face an obstacle, as discussing mental health is taboo. Mentally ill people are publicly humiliated in Pakistan, and sufferers are occasionally referred to as "pagal." Even parents ignore the symptoms of their children.

Review Article**Published Date:- 2022-05-31**[The surrey county lunatic asylum-an overview of some of the first admissions in 1863-1867](#)

In the 19th Century in much of Western Europe and North America the number and size of asylums increased hugely. In London, there was a wave of new asylums built in response to the 1808 County Asylums Act and the 1845 Lunacy Act, which required publicly funded care for those deemed mentally unwell. One such asylum was the Surrey County Lunatic Asylum which was built on the grounds which now house Springfield University Hospital in South West London.

This paper describes the admission records from Surrey County Lunatic Asylum, between 1863-1867, from information stored in the London Metropolitan Archives.

Although the terminology is different from that of today's, the picture the records paint is of an institution aiming at recovery rather than long-term incarceration which can be how asylums are now remembered. This more nuanced view is starting to be discussed more in public conversations about the topic. The optimism this may imbue is tempered by the shocking number of patients who died within the institution.

Case Report**Published Date:- 2022-05-31**[Fregoli syndrome in schizophrenia: about a case report](#)

A Fregoli syndrome, just like Capgras Syndrome, Double Subjective Syndrome, and Inter metamorphosis Syndrome, belongs to the group of delusion misidentification syndromes. It is a rare neuropsychiatric pathology that could affect the brain with repercussions on behavior. It is the belief that a familiar person is disguised as a strange person by taking a different physical appearance but remains the same person psychologically. This entity has heterogenous etiologies, occurring mainly in the setting of organic diseases, affective disorders, or schizophrenia. In this article, we proposed to expose a case of a schizophrenic patient who developed Fregoli syndrome.

[The role of cognitive distraction in sexual dysfunction in female medical resident doctors](#)

Introduction: Studies have shown a high prevalence of sexual dysfunction in Medical residents. Psychological, interpersonal, and sociocultural factors play an important role in sexual vulnerability. In addition, distraction or interference in the cognitive process of the erotic stimulus plays an important role in the development of sexual dysfunction.

The aim of our study was to determine the prevalence of sexual dysfunctions among a sample of married female medical resident doctors and to explore its relation with cognitive distraction during sex.

Subjects and methods: This was a cross-sectional study, which was carried out in October 2017 with 70 married female medical resident doctors. We used the Female Sexual Function Index (FSFI) for sexual functioning assessment and the cognitive distraction during sex scale.

Results: The mean age of women was 30.52 years. They had been married for 6.04 years on the mean. The majority (74%) had children. According to the FSFI, 40% of resident doctors had sexual dysfunction. The mean cognitive distraction during sex score was 4.37.

In our study, women with female sexual dysfunction reported significantly lower sexual desire score = 3.2 vs. 4.1 and $p = 0.03$) and arousal (score = 3.5 vs. 4.7 and $p = 10^{-3}$), decreased lubrication (score = 3.8 vs. 5.1; $p = 10^{-3}$), less sexual satisfaction (score = 3.9 vs. 5.8; $p = 10^{-3}$)

and higher orgasm difficulties (score = 3.4 vs. 5.1; $p = 10^{-3}$). Yet, our results highlighted that women with female sexual dysfunction presented significantly more cognitive distraction during sex (score = 3.9 vs. 4.6; $p = 0.03$).

Conclusion: According to our study, it appears that the prevalence of FSD in medical resident doctors is considerable. Besides, intrusive thoughts and cognitive distraction during sexual intercourse are importantly associated with sexual functioning. Therefore, several measures should be implemented in this population to prevent or act on these factors.

[The study of social and cultural values of Baiga tribes in the state of Madhya Pradesh, in India](#)

This paper is related to primary and secondary data collected from Anupur district, Madhya Pradesh. It is based on the social and cultural values of the Baiga tribes. Baiga tribe is the most backward tribe of Madhya Pradesh. These tribal people are found in this tribe are found in the following regions of Madhya Pradesh i.e. Eastern Satpura, Mandla, Dindori, Balaghat, Baghelkhand, Sidhi Janpads, Bilaspur, and even in the sarguju area of Chhatisgarh state. The tribe baiga has been defined as decedents from sorcerers or medicine men. The term baiga in the long past time was also applied in the central provinces to anyone who served as a village priest. The methodology used in this research paper is of descriptive and exploratory types. The objective of this paper is conducted through social and cultural values analysis sociological aspects with structure and functional approach.

[Organizational and methodological aspects of the work of a psychiatrist on a qualification category. Current status and prospects of psychiatric care in Russia](#)

Introduction: Preparing a report and passing certification for a qualification category is one of the forms of improving the professional level of each medical worker.

Material and methods: The analysis of 5 qualification works of a psychiatrist (1997, 2002, 2008, 2013, and 2018) was carried out for the mandatory requirements for the preparation of documents when they were submitted to the Attestation Commission on the assignment of a qualifying medical category in psychiatry and formation methodological approaches to their implementation. The depth of research was 27 years.

Results: Methodological approaches to the design of 11 mandatory sections provided for in the certification work of a psychiatrist presented for the award of a qualification category were considered.

Discussion: The attitude to the Administrative Regulations for the provision of state services for the assignment of a qualification category to specialists engaged in medical and pharmaceutical activities should be revised in terms of its synchronization with the official duties of a doctor.

Conclusion: The dynamic development of the regulatory, organizational, methodological, and scientific aspects of medicine and psychiatry, in particular, in the period of active digitalization of healthcare, requires periodic corporate discussion and correction of the Administrative Regulations for the provision of public services for assigning a qualification category to specialists engaged in medical and pharmaceutical activities.

Attention should be paid to the insufficient statistical processing of the submitted materials by applicants for the assignment of the 1st and highest categories, the lack of calculations, references, and comparisons of confidence intervals, and the reliability of the study.

Preparing a report and passing certification for a qualification category can be the first step in introducing a practitioner to scientific work through the generalization of personal work experience, worthy of speaking at a scientific and practical conference at the level of LU and the region and publishing abstracts in collections and scientific and practical journals.

The proposed methodological approaches are purely advisory in nature and can be used by psychiatrists when working on a qualification category at their discretion.
